

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (currently amended) A fermentation medium comprising a fermenting microorganism and a cytokinin-containing preparation that comprises a cytokinin at a concentration of at least 1 microMolar and effective to increase fermentation of the [a] microorganism in the fermentation medium.
2. (original) The fermentation medium of claim 1 wherein the cytokinin-containing preparation comprises a synthetic cytokinin having a purine heterocyclic base or a pyrimidine heterocyclic base.
3. (original) The fermentation medium of claim 2 wherein the purine heterocyclic base is an N⁶-substituted adenine or an optionally N⁶-substituted guanine.
4. (original) The fermentation medium of claim 2 wherein the synthetic cytokinin is selected from the group consisting of N⁶-benzyladenine, N⁶-benzyladenosine, N⁶-benzyladenine-3-glucoside, N⁶-benzyladenine-7-glucoside, N⁶-benzyladenine-9-glucoside, N⁶-benzyl-9-(2-tetrahydropyranyl)adenine, N⁶-benzyladenosine-5'-monophosphate, N⁶-gamma, gamma-dimethyl-allyl-aminopurine, dihydrozeatin, dihydrozeatin riboside, dihydrozeatin-7-beta-D-glucoside, dihydrozeatin-9-beta-D-glucoside, dihydrozeatin-O-glucoside, dihydrozeatin-O-glucoside riboside, dihydrozeatin riboside-5'-monophosphate, dihydrozeatin-O-acetyl; N⁶-isopentenyladenine, N⁶-isopentenyladenosine, N⁶-isopentenyladenosine-5'-monophosphate, N⁶-isopentenyladenine-7-glucoside, N⁶-isopentenyladenine-9-glucoside, 2-methylthio-N⁶-isopentenyladenosine, 2-methylthio-N⁶-isopentenyladenine, 2-thio-N⁶-isopentenyladenine, 2-benzylthio-N⁶-isopentenyladenine, 2-isopentenylamine, kinetin, kinetin riboside, kinetin-9-glucoside, kinetin riboside-5'-monophosphate, meta-topolin, meta-topolin riboside, meta-topolin-9-glucoside, ortho-topolin, ortho-topolin riboside, ortho-topolin-9-glucoside, trans-zeatin, trans-zeatin riboside, cis-zeatin, cis-zeatin riboside, trans-zeatin-7-glucoside, trans-zeatin-9-glucoside, trans-zeatin-O-glucoside, trans-zeatin-O-glucoside riboside, trans-zeatin

riboside-5'-monophosphate, trans-zeatin-O-acetyl, 2-chloro-trans-zeatin, 2-methylthio-trans-zeatin, and 2-methylthio-trans-zeatin riboside.

5. (currently amended) The fermentation medium of claim 4 wherein the cytokinin is present in the fermentation medium at a concentration of at least 5.0 [1.0] microM.
6. (original) The fermentation medium of claim 1 wherein the cytokinin-containing preparation comprises a plant extract.
7. (original) The fermentation medium of claim 6 wherein the plant is a member of the genus *Hordeum*.
8. (original) The fermentation medium of claim 1 wherein the cytokinin is present in the fermentation medium at a concentration effective to activate an AMP-activated protein kinase of the microorganism.
9. (original) The fermentation medium of claim 1 wherein the cytokinin is present in the fermentation medium at a concentration effective to increase uptake of a carbohydrate into the microorganism.
10. (original) The fermentation medium of claim 1 wherein the microorganism is a yeast and a member of the genus *Saccharomyces*.
11. (original) The fermentation medium of claim 1 wherein the fermentation medium is a liquid and wherein the fermentation comprises production of at least one of ethanol and carbon dioxide.
12. (original) The fermentation medium of claim 11 wherein the liquid is a beverage for human consumption.
13. (original) The fermentation medium of claim 1 wherein the fermentation medium is a dough and wherein the fermentation comprises production of carbon dioxide.
14. (currently amended) A method of increasing fermentation of a microorganism, comprising:

providing a cytokinin-containing preparation, and providing a fermentation medium; and combining the fermentation medium with the cytokinin-containing preparation, wherein the cytokinin-containing preparation is present in the fermentation medium at at least 1.0 microMolar to thereby ~~in an amount effective to~~ increase fermentation of a microorganism; and
inoculating the fermentation medium with a fermenting microorganism.

15. (original) The method of claim 14 wherein the cytokinin-containing preparation comprises a cytokinin that includes an N⁶-substituted adenine or an optionally N⁶-substituted guanine.
16. (original) The method of claim 14 wherein the cytokinin-containing preparation comprises a cytokinin selected from the group consisting of N⁶-benzyladenine, N⁶-benzyladenosine, N⁶-benzyladenine-3-glucoside, N⁶-benzyladenine-7-glucoside, N⁶-benzyladenine-9-glucoside, N⁶-benzyl-9-(2-tetrahydropyranyl)adenine, N⁶-benzyladenosine-5'-monophosphate, N⁶-gamma, gamma-dimethyl-allyl-aminopurine, dihydrozeatin, dihydrozeatin riboside, dihydrozeatin-7-beta-D-glucoside, dihydrozeatin-9-beta-D-glucoside, dihydrozeatin-O-glucoside, dihydrozeatin-O-glucoside riboside, dihydrozeatin riboside-5'-monophosphate, dihydrozeatin-O-acetyl; N⁶-isopentenyladenine, N⁶-isopentenyladenosine, N⁶-isopentenyladenosine-5'-monophosphate, N⁶-isopentenyladenine-7-glucoside, N⁶-isopentenyladenine-9-glucoside, 2-methylthio-N⁶-isopentenyladenosine, 2-methylthio-N⁶-isopentenyladenine, 2-thio-N⁶-isopentenyladenine, 2-benzylthio-N⁶-isopentenyladenine, 2-isopentenylamine, kinetin, kinetin riboside, kinetin-9-glucoside, kinetin riboside-5'-monophosphate, meta-topolin, meta-topolin riboside, meta-topolin-9-glucoside, ortho-topolin, ortho-topolin riboside, ortho-topolin-9-glucoside, trans-zeatin, trans-zeatin riboside, cis-zeatin, cis-zeatin riboside, trans-zeatin-7-glucoside, trans-zeatin-9-glucoside, trans-zeatin-O-glucoside, trans-zeatin-O-glucoside riboside, trans-zeatin riboside-5'-monophosphate, trans-zeatin-O-acetyl, 2-chloro-trans-zeatin, 2-methylthio-trans-zeatin, and 2-methylthio-trans-zeatin riboside.

17. (original) The method of claim 16 wherein the cytokinin is present in the fermentation medium at a concentration of at least 1.0 microM.
18. (original) The method of claim 14 wherein the cytokinin is present in the fermentation medium at a concentration effective to activate an AMP-activated protein kinase of the microorganism.
19. (original) The method of claim 14 wherein the cytokinin is present in the fermentation medium at a concentration effective to increase uptake of a carbohydrate into the microorganism.
20. (original) The fermentation medium of claim 1 wherein the microorganism is a yeast and a member of the genus *Saccharomyces*, wherein the fermentation medium is a liquid, and wherein the fermentation comprises production of at least one of ethanol and carbon dioxide.
21. (withdrawn) A method of marketing a product, comprising a step of providing information that a cytokinin increases fermentation of a microorganism.